



COLORECTAL

Did you know?

- Neuroendocrine Tumours (NETs) are the second most common cancer of the digestive tract³
- NETs of the gastrointestinal tract are often found in the appendix, colon and rectum¹⁰
- Colon NETs do not exhibit symptoms until late into the progression of the disease, often at a point where the cancer has spread. Common symptoms of colon NETs include diarrhea, abdominal pain and weight loss¹⁰
- Rectal NETs are often found unexpectedly during colorectal cancer screening. Common symptoms of rectal NETs include discomfort, pain and changes in bowel habits¹⁰
- NETs of the appendix are often found after examination and surgery for appendicitis¹⁰

When it comes to NETs,
AWARENESS OF THIS DISEASE IS KEY TO DIAGNOSING IT.

If healthcare professionals do not think about NETs, they will not look for them.⁹

Help us raise awareness of NETs, including NETs of the digestive tract.

LEARN MORE ABOUT DIAGNOSING NETS



CCRAN
Colorectal Cancer
Resource &
Action Network



Neuroendocrine (NET) Cancer Day is an annual **event held on the 10th of November**, created to increase awareness of Neuroendocrine (NET) cancers and to provide a voice to the NET community for improved diagnostics, treatments, information, care and research.

NETs are a unique group of cancers that arise from neuroendocrine cells.¹ These tumours can arise in many organs of the body.²

NETs affect thousands of people living in Canada every year, and this number has been increasing.²

NETs generally affect people over the age of 50.³

Symptoms of NETs vary and are often non-specific.¹ This is due to their ability to secrete an excess of different hormones.⁴

The symptoms may depend on the organ where the NET is located, and can include abdominal pain, skin flushing, diarrhea, cough, and back pain.⁵

As a distinct class of tumours, NETs require specialized tests and treatments.¹


As with other types of cancer, early diagnosis and treatment can improve survival.^{6,7}

Unfortunately, most NETs are not found until it is too late. Results from a global survey show an average delay of 4 years from first symptoms to diagnosis.⁸

This is generally due to the perception that NETs are uncommon and a lack of awareness of the disease.^{1,9}

World NET Cancer Day is about drawing attention to this uncommon and often misdiagnosed form of cancer and starting a global conversation amongst communities, medical professionals and governments.

#LetsTalkAboutNETs



1. Raphael MJ, Chan DL, Law C, Singh S. Principles of diagnosis and management of neuroendocrine tumours. *CMAJ*. 2017;189(10):E398-E404. doi:10.1503/cmaj.160771

2. Hallet J, Law CHL, Cukier M, Saskin R, Liu N, Singh S. Exploring the rising incidence of neuroendocrine tumors: A population-based analysis of epidemiology, metastatic presentation, and outcomes. *Cancer*. 2015;121(4):589-597. doi:10.1002/cncr.29099

3. Yao JC, Hassan M, Phan A, et al. One hundred years after "carcinoid": epidemiology of and prognostic factors for neuroendocrine tumors in 35,825 cases in the United States. *J Clin Oncol*. 2008;26(18):3063-3072. doi:10.1200/JCO.2007.15.4377

4. Neuroendocrine tumors - Symptoms and causes - Mayo Clinic. <https://www.mayoclinic.org/diseases-conditions/neuroendocrine-tumors/symptoms-causes/syc-20354132>. Accessed October 25, 2021.

5. Ter-Minassian M, Chan JA, Hooshmand SM, et al. Clinical presentation, recurrence, and survival in patients with neuroendocrine tumors: results from a prospective institutional database. *Endocr Relat Cancer*. 2013;20(2):187-196. doi:10.1530/ERC-12-0340

6. Ahmed A, Turner G, King B, et al. Midgut neuroendocrine tumours with liver metastases: results of the UKINETS study. *Endocr Relat Cancer*. 2009;16(3):885.

7. Hill JS, McPhee JT, McDade TP, et al. Pancreatic neuroendocrine tumors: the impact of surgical resection on survival. *Cancer Interdiscip Int J Am Cancer Soc*. 2009;115(4):741-751.

8. Singh S, Granberg D, Wolin E, et al. Patient-Reported Burden of a Neuroendocrine Tumor (NET) Diagnosis: Results From the First Global Survey of Patients With NETs. *J Glob Oncol*. 2016;3(1):43-53. doi:10.1200/JGO.2015.002980

9. Aggarwal G, Obideen K, Wehbi M. Carcinoid tumors: what should increase our suspicion? *Cleve Clin J Med*. 2008;75(12):849-855. doi:10.3949/ccjm.75a.08002

10. Shah R, Nalamati SPM. Neuroendocrine tumors of the colon and rectum. *Semin Colon Rectal Surg*. 2015;26(2):60-63. doi:https://doi.org/10.1053/j.scrs.2015.01.002